

Universality in heavy-fermion systems with general degeneracy

N. Tsujii¹, H. Kontani², and K. Yoshimura³

¹*National Institute for Materials Science, Sengen 1-2-1, Tsukuba, 305-0047, Japan*

²*Graduate School of Science, Nagoya University,*

Fro-cho, Chikusa-ku, Nagoya City 464-8602, Japan and

³*Graduate School of Science, Kyoto University, Kyoto 606-8502, Japan*

(Dated: February 2, 2008)

We discuss the relation between the T^2 -coefficient of electrical resistivity A and the T -linear specific-heat coefficient γ for heavy-fermion systems with general N , where N is the degeneracy of quasi-particles. A set of experimental data reveals that the Kadowaki-Woods relation; $A/\gamma^2 = 1 \times 10^{-5} \mu\Omega(\text{K mol/mJ})^2$, collapses remarkably for large- N systems, although this relation has been regarded to be commonly applicable to the Fermi-liquids. Instead, based on the Fermi-liquid theory we propose a new relation; $\tilde{A}/\tilde{\gamma}^2 = 1 \times 10^{-5}$ with $\tilde{A} = A/\frac{1}{2}N(N-1)$ and $\tilde{\gamma} = \gamma/\frac{1}{2}N(N-1)$. This new relation exhibits an excellent agreement with the data for whole the range of degenerate heavy-fermions.

PACS numbers: 71.10.Ay, 71.27.+a, 75.30.Mb

The Fermi-liquid theory [1] is the most fundamental one to understand the electronic state of metallic systems. This theory has achieved a great success in describing not only the electronic properties of normal metals but also unusual properties of the strongly-correlated electron systems [2, 3] like f -electron based heavy-fermion compounds [2, 3, 4, 5] and d -electron based intermetallics and oxides [2, 6]. In this theory, the effect of electron-electron interactions are involved in the effective mass of quasi-particles, m^* . This enables a very simple representation of physical properties: the electronic specific heat C and the electrical resistivity ρ are described as $C = \gamma T$ and $\rho = AT^2$ with $\gamma \propto m^*$ and $A \propto m^{*2}$. Such a temperature dependence has been actually observed in numerous kind of metals. Moreover, this description implies that the ratio A/γ^2 does not depend on m^* , resulting in the universal value of A/γ^2 . In fact, it has been revealed that many f -electron based systems show the universal behavior; $A/\gamma^2 = 1.0 \times 10^{-5} \mu\Omega\text{cm}(\text{Kmol/mJ})^2$ [7], called as the Kadowaki-Woods (KW) relation. The KW-relation has therefore been accepted as the most essential relation showing the validity of the Fermi-liquid theory.

Recently, however, significant and systematic deviations from the relation have been observed in many heavy-fermion compounds, in spite that they apparently show Fermi-liquid behavior at low temperature [8]. This class of compounds includes Yb-based compounds like YbCu₅, YbAgCu₄, YbCuAl, YbNi₂Ge₂, YbInCu₄, YbAl₃, and Ce-based compounds like CeNi₉Si₄ [9] and CeSn₃. Notably, the deviations in these systems are almost 'universal'; $A/\gamma^2 \approx 0.4 \times 10^{-6} \mu\Omega\text{cm}(\text{Kmol/mJ})^2$. This systematic and large deviation cannot be explained by specific characters of materials, like carrier density, band structure, anisotropy, etc. Instead, there seems to exist a common physical origin. The origin of this deviation therefore rises an important issue for the generality of the Fermi-liquid theory.

Very recently, a theoretical work based on the Fermi-

liquid theory has suggested [10] that the values of A/γ^2 , so far considered to be unique and independent on materials, do depend on the number of degeneracy of quasi-particles N . For isolated atoms, N is defined as $N = 2J + 1$ with J the total angular momentum. In solids, N can vary due to the competition between the crystal-field splitting Δ and the Kondo temperature T_K . For $T_K < \Delta$, the low-temperature properties are basically explained by $N = 2$ ($S = 1/2$) Kondo model, since most of the degeneracy are lost due to the large Δ [5, 11, 12]. For $T_K > \Delta$, on the contrary, the crystal field splittings are covered by the large Kondo effect, and the degeneracy are almost preserved down to low temperatures. In this case, the theory [10] gives the failed universality of A/γ^2 .

In this paper, we make a quantitative comparison of the experimental data in ref.[8] and several recent works with the theoretical results of ref.[10]. The results display a beautiful agreement between experiments and theory. Furthermore, we propose an advanced relation for A and γ based on these results. Using \tilde{A} and $\tilde{\gamma}$, the values of A and γ normalized by $\frac{1}{2}N(N-1)$, we show that these two values of heavy-fermion systems with general N are related by a very simple formula; $\tilde{A}/\tilde{\gamma}^2 = 1 \times 10^{-5} \mu\Omega\text{cm}(\text{Kmol/mJ})^2$. This new relation, namely, the 'grand-KW-relation', will be an important waymark for the research of strongly-correlated electron systems with degeneracies, and remarkably extends the validity of the Fermi-liquid theory.

At first, we briefly describe the theoretical results of ref. [10]. For the case of strong-coupling limit where $m^*/m \gg 1$ (m^* and m being the mass of heavy quasi-particles and free electrons, respectively), the orbitally-degenerate periodic Anderson (ODPA) model gives [13]:

$$A = \frac{\hbar k_B^2}{e^2} \frac{3\pi^6}{2k_F^4 a^3} N(N-1) \Gamma_{\text{loc}}^2(0,0) \rho_f^4(0), \quad (1)$$

$$\gamma = N_A k_B^2 \frac{\pi^2}{6} N(N-1) \Gamma_{\text{loc}}(0,0) \rho_f^2(0). \quad (2)$$




FIG. 2: The plot of \tilde{A} and $\tilde{\gamma}$ of heavy-fermion systems. \tilde{A} and $\tilde{\gamma}$ are the divided values of A and γ by $\frac{1}{2}N(N-1)$, respectively. N of the U-based compounds is tentatively assumed to be 2. The dotted line represents the grand KW-relation (4) given in the text; $\tilde{A}/\tilde{\gamma}^2 = 1 \times 10^{-5} \mu\Omega(\text{K mol/mJ})^2$.

the value of $\tilde{A}/\tilde{\gamma}^2$ defined at the low temperature limit ($T \rightarrow 0$), follows the relation (4) even in the vicinity of QCP. This is because our theory is derived for the limit of $T \rightarrow 0$. It should also be noticeable to point that the theoretical calculations of A/γ^2 for $N = 2$ based on the spin-fluctuation theory show that the ratio is approximately independent of the distance from the QCP [20, 21]. In fact, A/γ^2 of YbRh₂Si₂ and CeInCo₅, both of which are considered to be in the vicinity of QCP, are almost a constant as external magnetic field is varied [22, 23]. Meanwhile, for the case of ‘very’ close to the QCP, where T_{coh} is quite low, a deviation from the universality may be observed, as is suggested theoretically [20] and experimentally on YbRh₂(Si,Ge)₂ [24]. This deviation however seems to occur in an extremely narrow condition where the Fermi-liquid description is probably not valid. Except for such extreme cases, the grand-KW relation is one of the common behavior of Fermi liquids, even in systems close to the QCP. Note that the f -orbital degeneracy will

stabilize the Fermi-liquid state, because N -dependence of T_K ($\propto e^{-1/\rho^N J_K}$) will be much prominent than that of T_N ($\propto N^2 J_{\text{RKKY}}$). A large mass-enhancement is realized with relatively higher T_K when $N > 2$.

There should, of course, exist exceptions. As is seen in the formula (3), the ratio A/γ^2 as well as that of $\tilde{A}/\tilde{\gamma}^2$ depend on the carrier concentration n , wave number at the Fermi energy k_F , and so on. If one of these values are extremely different from typical ones, $\tilde{A}/\tilde{\gamma}^2$ can deviate remarkably. Such an example is CePd₃. In Fig. 2, one can see that CePd₃ shows a large deviation from eq. (4), though CePd₃ well agrees with the original KW-relation [7]. This discrepancy results from the large degeneracy, $N = 6$ for CePd₃ [25]. It should be noted that CePd₃ has very small carrier-concentration (0.3 electrons per f.u.) [26]. The A/γ^2 value is found to depend on n as proportional to $n^{-4/3}$ from eq. (3) and also from other theoretical studies [2, 20, 21, 27]. Taking this into consideration, the deviation of CePd₃ from the universal line is reasonable. Similar deviation is reported for the Kondo semiconductor CeNiSn [28]. Anomalously large A value (54 $\mu\Omega\text{cm}/\text{K}^2$) compared to its γ (40 mJ/mol K²) has been attributed to its extremely low carrier concentration [28].

The compounds CeNi ($N = 6$) and YbCuAl ($N = 8$) also show slight deviations, possibly due to the error in the N estimations. For other exceptions, YbInAu₂ and Yb₂Co₃Ga₉, we have no explanation for the origin of deviation. Other causes such as multi-Fermi-surface effect [9] may have to be considered. In addition, strong anisotropy of the Fermi surface can cause deviation from the universal relation [2, 29]. This effect would be in general more prominent in d -electron systems [30, 31].

For U-based compounds, its degeneracy has been the subject of arguments. If the $5f$ -electrons are well localized, N can be determined experimentally, as in the case of UPd₃ [32]. In most of U-compounds, however, it is considered that the $5f$ -electrons have more-itinerant character than $4f$, since $5f$ -orbitals are spatially more expanded. The definition of N in U-compounds is therefore ambiguous. Here, one can see in Fig. 1 and Fig. 2 that those U-compounds well agree with the theoretical prediction for $N = 2$. This can lead us to the possibility that the orbital degree of freedom is quenched and only the spin degree of freedom participates in the Fermi-liquid state in these $5f$ -systems, similar to transition metals. Although the estimation of N from the A/γ^2 plot is not conclusive, this plot may serve as a hint to discuss the puzzling $5f$ -electrons.

In addition, we note that the grand-KW-relation is also powerful to describe the pressure dependent properties of heavy-fermion systems. In CeCu₂Ge₂ (or YbNi₂Ge₂), it is suggested that the value of A/γ^2 reduces (or increases) about 25 times at high pressures [33, 34] probably due to the change of N by pressures. In our plot of \tilde{A} and $\tilde{\gamma}$, these crossover would be described on the single scaling without breaking the universality. This situation may be hence ideal for the continuity principle of the Landau

Fermi-liquid theory [35]. Current interests in strongly-correlated electron systems are extended to the orbitally-degenerate cases. Hence, the grand-KW-relation will be one of the most fundamental relation in Fermi-liquid systems. We also comment that the effect of the degeneracy must be taken into consideration in many other physical

quanta, like anomalous Hall effect [36], etc. The analysis using the ODP model will be henceforth indispensable.

Authors acknowledge K. Yamada, A. Mitsuda, Y. Aoki, G. Kido and H. Kitazawa for fruitful discussions and comments.

-
- [1] L. D. Landau, Sov. Phys. JETP **3**, 920 (1956); *ibid.* **8**, 70 (1959).
 - [2] K. Yamada, *Electron Correlation in Metals*, Cambridge Univ. Press, US, 2004.
 - [3] Y. Yanase, T. Jujo, T. Nomura, H. Ikeda, T. Hotta, and K. Yamada, Phys. Rep. **387**, 1 (2003).
 - [4] For a review, see G. R. Stewart, Rev. Mod. Phys. **56**, 755 (1984), C. M. Varma, Comments Solid State Phys. **11**, 221 (1985), etc.
 - [5] N. B. Brandt and V.V. Moshchalkov, Adv. Phys. **33**, 373 (1984).
 - [6] T. Moriya and K. Ueda, Rep. Prog. Phys. **66**, 1299 (2003).
 - [7] K. Kadowaki and S. B. Woods, Solid State Commun. **58**, 507 (1986).
 - [8] N. Tsujii, K. Yoshimura and K. Kosuge, J. Phys.: Condens. Matter. **15**, 1993 (2003).
 - [9] H. Michor, St. Berger, M. El-Hagary, C. Paul, E. Bauer, G. Hilscher, P. Rogl and G. Giester, Phys. Rev. B **67**, 224428 (2003).
 - [10] H. Kontani, J. Phys. Soc. Jpn. **73**, 515 (2004).
 - [11] M. J. Besnus, J. P. Kappler, P. Lehmann and A. Meyer, Solid State Commun. **55**, 779 (1985).
 - [12] T. Fujita, K. Satoh, Y. Onuki and T. Komatsubara, J. Mag. Mag. Mater. **47-48**, 66 (1985).
 - [13] The eq.(1) and (2) explicitly describes the units like k_B and \hbar , that were omitted in the equations of ref. [10].
 - [14] Here the free electron model $k_F = (3\pi^2 n)^{1/3}$ (n being the carrier concentration) is employed. For numerical estimation, we have assumed $n^{4/3} a^3 \approx 1 \times 10^8 \text{ cm}^{-1}$. This assumption corresponds to 3.3 carriers per formula unit for $a = 5\text{\AA}$ or to 2.8 carriers per f.u. for $a = 4\text{\AA}$, which seems to be typical values for them. Formula unit is to include only one rare-earth ion.
 - [15] For the values of N , we have employed the data listed in ref. [8], most of which were determined by fitting the specific heat using the impurity-Kondo models. Although the real system is periodic, we believe that this estimation of N is reliable, because these fittings are performed for the data at relatively high temperatures, at which intersite effect are not prominent. However, it is difficult to determine the value of N explicitly when Δ (crystal-field splitting) and T_K are comparable.
 - [16] A. Mitsuda *et al.*, in preparation.
 - [17] N. Takeda and M. Ishikawa, J. Phys.: Condens. Matter. **15**, L229 (2003).
 - [18] S. Sanada, Y. Aoki, D. Kikuchi, H. Sugawara, H. Sato, J. Phys. Soc. Jpn., to be published.
 - [19] The lowest crystal-field level in CeB₆ is considered to be a quartet. However, we note that the Fermi-liquid behavior in this system is only observed below 0.5 K, which is much lower than the quadrupolar-ordering temperature of 3.3 K [N. Sato, A. Sumiyama, S. Kunii, H. Nagano, and T. Kasuya, J. Phys. Soc. Jpn. **54**, 1923 (1985)]. Therefore, the degeneracy relevant to the Fermi-liquid behavior should be much smaller than 4.
 - [20] T. Takimoto and T. Moriya, Solid State Commun. **99**, 457 (1996).
 - [21] M.A. Continentino, Eur. Phys. J. B **13**, 31 (2000).
 - [22] P. Gegenwart, J. Custers, C. Geibel, K. Neumaier, T. Tayama, K. Tenya, O. Trovarelli and F. Steglich, Phys. Rev. Lett. **89**, 056402 (2002).
 - [23] A. Bianchi, R. Movshovich, I. Vekhter, P.G. Pagliuso and J.L. Sarrao, Phys. Rev. Lett. **91**, 257001 (2003).
 - [24] J. Custers, P. Gegenwart, H. Wilhelm, K. Neumaier, Y. Yokiwa, O. Trovarelli, C. Geibel, F. Steglich, C. Pépin and P. Coleman, Nature **424**, 524 (2003).
 - [25] A. P. Murani, R. Raphael, Z. A. Bowden and R. S. Eccleston, Phys. Rev. B **53**, 8188 (1996).
 - [26] B. C. Webb, A. J. Sievers and T. Mihalisin, Phys. Rev. Lett. **57**, 1951 (1986).
 - [27] In the reference [21], $A/\gamma^2 \propto \rho_0 \propto (N_c^2 \tau_{Fc})^{-1} \propto k_F^2/n^2$. Assuming $k_F = (3\pi^2 n)^{1/3}$, this gives the $n^{-4/3}$ dependence.
 - [28] T. Terashima, C. Terakura, S. Uji, H. Aoki, Y. Echizen and T. Takabatake, Phys. Rev. B **66**, 075127 (2002).
 - [29] H. Fukazawa *et al.*, unpublished.
 - [30] N.E. Hussey, cond-mat/0409252, (2004).
 - [31] A/γ^2 of transition metals like Pd and Pt also deviates from the KW-relation, and is the same order as that of the degenerate heavy-fermion systems ($N \approx 8$); $A/\gamma^2 = 0.4 \times 10^{-6} \mu\Omega\text{cm}(\text{K mol/mJ})^2$ [M. J. Rice, Phys. Rev. Lett. **20**, 1439 (1968)]. However, these transition metals are not likely to hold the universal relation, eq.(4), because N of these metals would be close to 2 due to the quenching of the orbital moment. A/γ^2 of these metals is hence much smaller than eq.(4). Here, note that the theory [10] we use in the present work is based on the strong-coupling case ($m^*/m \gg 1$), which is apparently not the case of these metals. For such weak-coupling cases, deviation from the KW-relation is suggested theoretically in [K. Miyake, T. Matsuura and C. M. Varma, Solid State Commun. **71**, 1149 (1989)].
 - [32] W. J. L. Buyers, A. F. Murray, T. M. Holden, E. C. Svensson, P. de V. DuPlessis, G. H. Lander and O. Vogt, Physica B+C **102**, 291 (1980).
 - [33] D. Jaccard, H. Wilhelm, K. Alami-Yadri and E. Vargoz, Physica B **259-261**, 1(1999).
 - [34] G. Knebel, D. Braithwaite, G. Lapertot, P. C. Canfield and J. Flouquet, J. Phys.: Condens. Matter **13**, 10935 (2001).
 - [35] P. W. Anderson, *Basic Notions of Condensed Matter Physics*, Addison-Wesley Publishing Company Inc., 1984.
 - [36] H. Kontani and K. Yamada, J. Phys. Soc. Jpn. **63**, 2627 (1994).